

PROJECTIVE STRUCTURES, GRAFTING, AND MEASURED LAMINATIONS

DAVID DUMAS AND MICHAEL WOLF

ABSTRACT. We show that grafting any fixed hyperbolic surface defines a homeomorphism from the space of measured laminations to Teichmüller space, complementing a result of Scannell-Wolf on grafting by a fixed lamination. This result is used to study the relationship between the complex-analytic and geometric coordinate systems for the space of complex projective ($\mathbb{CP}^1$) structures on a surface.

We also study the rays in Teichmüller space associated to the grafting coordinates, obtaining estimates for extremal and hyperbolic length functions and their derivatives along these grafting rays.

1. INTRODUCTION

In this paper we compare two perspectives on the theory of complex projective structures on surfaces by studying the grafting map of a hyperbolic surface.

A complex projective (or $\mathbb{CP}^1$) structure on a compact surface S is an atlas of charts with values in $\mathbb{CP}^1$ and Möbius transition functions. Let $\mathcal{P}(S)$ denote the space of (isotopy classes of) marked complex projective structures on S , and let $\mathcal{T}(S)$ be the Teichmüller space of (isotopy classes of) marked complex structures on S . Because Möbius maps are holomorphic, there is a forgetful projection $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$.

An analytic tradition, having much in common with univalent function theory, parameterizes the fiber $\pi^{-1}(X)$ using the Schwarzian derivative, identifying $\mathcal{P}(S)$ with the total space of the bundle $\mathcal{Q}(S) \rightarrow \mathcal{T}(S)$ of holomorphic quadratic differentials.

A second, more synthetic geometric description of $\mathcal{P}(S)$ is due to Thurston, and proceeds through the operation of *grafting* – a construction which traces its roots back at least to Klein [Kle, §50, p. 230], with a modern history developed by many authors ([Mas1], [Hej], [ST], [Gol], [GKM], [Tan], [McM], [SW]). The simplest example of grafting may be described as follows.

Start with a hyperbolic surface $X \in \mathcal{T}(S)$ and a simple closed geodesic γ on X ; then construct a new surface by removing γ from X and replacing it

Date: November 30, 2007.

2000 Mathematics Subject Classification. Primary 30F60.

The first author was partially supported by an NSF postdoctoral research fellowship.

The second author was partially supported by NSF grants DMS-0139877 and DMS-0505603.

with the Euclidean cylinder $\gamma \times [0, t]$. The result is $\text{Gr}_{t\gamma} X$, the *grafting of X by $t\gamma$* , which is a surface with a $(C^{1,1}$ Riemannian) metric composed of alternately flat or hyperbolic pieces. Furthermore, $\text{Gr}_{t\gamma} X$ has a canonical projective structure that combines the Fuchsian uniformization of X and the Euclidean structure of the cylinder $\gamma \times [0, t]$ (for details, see [SW, §1] [Tan, §2] [KT]).

Thurston showed that grafting extends naturally from weighted simple closed geodesics to the space $\mathcal{ML}(S)$ of measured geodesic laminations, and thus defines a map

$$(1.1) \quad \text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{P}(S).$$

Moreover, this map is a homeomorphism; for a proof of this result, see [KT].

A natural problem is to relate the analytic and geometric perspectives on the space of projective structures, for example by comparing the product structure of $\mathcal{ML}(S) \times \mathcal{T}(S) \simeq \mathcal{P}(S)$ to the bundle structure induced by the projection $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$.

Results on grafting. We compare these two perspectives by studying the *conformal grafting map* $\text{gr} = \pi \circ \text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{T}(S)$, i.e. $\text{gr}_\lambda X$ is the conformal structure which underlies the projective structure $\text{Gr}_\lambda X$. Fixing either of the two coordinates we have the X -grafting map $\text{gr}_\bullet X : \mathcal{ML}(S) \rightarrow \mathcal{T}(S)$ and the λ -grafting map $\text{gr}_\lambda : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$. These maps reflect how the base coordinate of the complex-analytic fibration $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ is related to the geometric product structure $\mathcal{ML}(S) \times \mathcal{T}(S) \simeq \mathcal{P}(S)$. Our main result is

Theorem 1.1. *For each $X \in \mathcal{T}(S)$, the X -grafting map $\text{gr}_\bullet X : \mathcal{ML}(S) \rightarrow \mathcal{T}(S)$ is a bitangentiable homeomorphism.*

Momentarily deferring a brief discussion of the term *bitangentiable homeomorphism*, we note that this theorem is a natural complement to the result of Scannell-Wolf on the λ -grafting map.

Theorem 1.2 (Scannell-Wolf [SW, Thm. A]). *For each $\lambda \in \mathcal{ML}(S)$, the map $\text{gr}_\lambda : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is a real-analytic diffeomorphism.*

The discrepancy between diffeomorphism and bitangentiable homeomorphism in Theorems 1.1 and 1.2 is related to the lack of a natural differentiable structure on $\mathcal{ML}(S)$. Bonahon showed that grafting is differentiable in the weak sense of being *tangentiable*; see §2 below or [Bon5] for details.

Returning to the original problem of comparing different coordinate systems for $\mathcal{P}(S)$, Theorems 1.1 and 1.2 can be used to study the fiber $P(X) = \pi^{-1}(X)$ and its relation to the grafting coordinates. Let us denote the two factors of the map $\text{Gr}^{-1} : \mathcal{P}(S) \rightarrow \mathcal{ML}(S) \times \mathcal{T}(S)$ by

$$\text{Gr}^{-1}(Z) = (p_{\mathcal{ML}}(Z), p_{\mathcal{T}}(Z)).$$




FIGURE 1. The bundle $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ of $\mathbb{CP}^1$ structures over Teichmüller space and the product structure $\text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \simeq \mathcal{P}(S)$ induced by grafting.

Thus the maps $p_{\mathcal{ML}} : \mathcal{P}(S) \rightarrow \mathcal{ML}(S)$ and $p_{\mathcal{T}} : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ send a projective structure to one of its two grafting coordinates, and we think of them as projections. We prove:

Corollary 1.3. *For each $X \in \mathcal{T}(S)$ the restriction $p_{\mathcal{ML}}|_{P(X)} : P(X) \rightarrow \mathcal{ML}(S)$ is a bitangentiable homeomorphism, and $p_{\mathcal{T}}|_{P(X)} : P(X) \rightarrow \mathcal{T}(S)$ is a C^1 diffeomorphism.*

This corollary improves the existing regularity results for these projection maps, from which it was known that that $p_{\mathcal{ML}}|_{P(X)}$ is a homeomorphism (a corollary of Theorem 1.2, see [D2, §4]) and that $p_{\mathcal{T}}|_{P(X)}$ is a proper C^1 map of degree 1 (see [Bon5, Thm. 3] and [D1, Lem. 7.6, Thm. 1.1]). The relationship between the maps $p_{\mathcal{T}}$, $p_{\mathcal{ML}}$, and π is represented schematically in Figure 1.

One can also apply Theorem 1.1 and Corollary 1.3 to study the *pruning map* (the inverse of grafting) and the parameterization of quasi-Fuchsian manifolds by their convex hull geometry. We explore these directions in §5.

Methods. From a general perspective, the proof of the main theorem relies on relating two established techniques in hyperbolic geometry. The first is the analytic study of the prescribed curvature (Liouville) equation (see e.g. [Wol4] [ZT], [Tro]), and the second is the complex duality between bending

and twisting (see e.g. [Wol1], [Wol2], [Wol3], [Pla], [Ser1], [Ser2]) and its generalization to complex earthquakes as a duality between grafting and shearing (see [Bon5], [McM], [EMM]).

The first strand occurs in the proof in [SW] of Theorem 1.2, where standard geometric analytic techniques were applied to the curvature equation to understand how the λ -grafting map changed under small perturbations of the Riemann surface. Such techniques might be applicable to the analogous problem (of the main Theorem 1.1) of understanding how the X -grafting map varies under small perturbations of the measured lamination, but it would necessarily be more involved, due to the local structure of the space $\mathcal{ML}(S)$ being more complicated than that of the space $\mathcal{T}(S)$.

Fortunately, most of the required details for this study of $\mathcal{ML}(S)$ are already in the literature: here we make heavy use of Bonahon's work (following Thurston [Thu3]) on the deformation theory of $\mathcal{ML}(S)$ (see [Bon2] [Bon3] [Bon4] [Bon5]). In particular, the crux of our proof relies on Bonahon's observation that there is a sense in which infinitesimal grafting is complex linear. This complex analyticity, in keeping with the second tradition discussed above, implies that a study of the effect on grafting of infinitesimally changing the measured lamination is, by duality, a study of the effect on grafting of infinitesimally shearing the hyperbolic surface. Thus we may apply the analysis in the proof of Theorem 1.2 to the problem of the main Theorem 1.1.

Grafting rays. In a final section, we study the coordinate system on Teichmüller space induced by the X -grafting homeomorphism $\mathcal{ML}(S) \rightarrow \mathcal{T}(S)$, and analyze the behavior of extremal and hyperbolic length functions on *grafting rays*—paths in $\mathcal{T}(S)$ of the form $t \mapsto \text{gr}_{t\lambda} X$. We show:

Theorem 1.4. *For each $X \in \mathcal{T}(S)$ and $\lambda \in \mathcal{ML}(S)$ the extremal length of λ on $\text{gr}_{t\lambda} X$ is monotone decreasing for all $t \gg 0$ and is asymptotic to $\frac{\ell(\lambda, X)}{t}$, where $\ell(\lambda, X)$ is the hyperbolic length of λ on X .*

Theorem 1.5. *For each $X \in \mathcal{T}(S)$ and any simple closed hyperbolic geodesic $\gamma \in \mathcal{ML}(S)$, the hyperbolic length of γ on $\text{gr}_{t\gamma} X$ is monotone decreasing for all $t \gg 0$ and is asymptotic to $\pi \frac{\ell(\lambda, X)}{t}$.*

The monotonicity and asymptotic behavior described in Theorems 1.4 and 1.5 are combined with explicit estimates on the derivatives of length functions in Theorems 6.2 and 6.6 below.

Organization of the paper.

§2 presents the infinitesimal version of the main theorem (Theorem 2.6), after introducing the necessary background on measured laminations and grafting. The reduction to an infinitesimal statement is modeled on the argument of Scannell-Wolf in [SW], and uses Tanigawa's properness theorem for grafting (Theorem 2.1).

§3 describes shearing deformations of hyperbolic surfaces, closely following the work of Bonahon on shearing coordinates for Teichmüller space. Portions of this section are more expository than is strictly necessary for the proofs of the of the results, but we feel they help make the arguments easier to understand. The discussion culminates with the crucial complex-linearity result of Bonahon (Theorem 3.4) that is used in the proof of Theorem 2.6.

§4 is devoted to the proofs of Theorems 2.6 and 1.1, which follow using the theory developed in §§2-3.

§5 collects some applications of the main theorem, including the proof of Corollary 1.3 and a rigidity result for quasi-Fuchsian manifolds.

§6 discusses the grafting coordinates for Teichmüller space and the asymptotic behavior of extremal and hyperbolic length functions on grafting rays; Theorems 1.4 and 1.5 and associated derivative estimates are proved here.

Acknowledgements. The authors thank Francis Bonahon, Howard Masur, Yair Minsky, and Robert Penner for stimulating discussions related to this work. They also thank the referee for several suggestions that improved the paper.

2. GRAFTING AND INFINITESIMAL GRAFTING

We begin with some background on measured laminations, grafting, and tangentiability, which are needed to formulate the main technical result (Theorem 2.6).

Laminations. As in the introduction, S denotes a compact smooth surface of genus $g > 1$ and $\mathcal{T}(S)$ is the Teichmüller space of marked hyperbolic (or conformal) structures on S . We often use $X \in \mathcal{T}(S)$ to represent a particular hyperbolic surface in a given marked equivalence class.

Let $\mathcal{S}$ denote the set of free homotopy classes of simple closed curves on S ; we implicitly identify $\gamma \in \mathcal{S}$ with its geodesic representative on a hyperbolic surface $X \in \mathcal{T}(S)$.

A *geodesic lamination* Λ on a hyperbolic surface X is a foliation of a closed subset of X by complete, simple geodesics. Examples of geodesic laminations include simple closed hyperbolic geodesics $\gamma \in \mathcal{S}$ and disjoint unions thereof.

The notion of a geodesic lamination is actually independent of the particular choice of X , in that a geodesic lamination on X determines a geodesic lamination for any other hyperbolic structure $Y \in \mathcal{T}(S)$ in a canonical way (see for example [Bon3, §1]). Thus we speak of a geodesic lamination on S , suppressing the choice of a particular metric. Let $\mathcal{GL}(S)$ denote the set of all geodesic laminations on S with the topology of Hausdorff convergence of closed sets.

A geodesic lamination $\Lambda \in \mathcal{GL}(S)$ is *maximal* if it is not properly contained in another geodesic lamination, in which case the complement of Λ

in S is a union of ideal triangles. Every geodesic lamination is contained in a maximal one, though not necessarily uniquely.

Measured laminations. A *transverse measure* μ on a geodesic lamination Λ is an assignment of a positive Borel measure to each compact transversal to Λ in a manner compatible with splitting and isotopy of transversals. Such a measure μ has *full support* if there is no proper sublamination $\Lambda' \subset \Lambda$ such that μ assigns the zero measure to transversals disjoint from Λ' .

Let $\mathcal{ML}(S)$ denote the space of *measured geodesic laminations* on S , i.e. pairs $\lambda = (\Lambda, \mu)$ where $\Lambda \in \mathcal{GL}(S)$ and μ is a transverse measure on Λ of full support. We denote by $\lambda(\tau)$ the total measure assigned to a transversal τ by $\lambda \in \mathcal{ML}(S)$.

The topology on $\mathcal{ML}(S)$ is that of weak-* convergence of measures on compact transversals. The underlying geodesic lamination of $\lambda \in \mathcal{ML}(S)$ is the *support* of λ , written $\text{supp}(\lambda) \in \mathcal{GL}(S)$. The space $\mathcal{ML}(S)$ has an action of $\mathbb{R}^+$ by multiplication of transverse measures; the empty lamination $0 \in \mathcal{ML}(S)$ is the unique fixed point of this action.

For any simple closed geodesic $\gamma \in \mathcal{S}$, there is a measured geodesic lamination (also γ) that assigns to a transversal τ the counting measure on $\tau \cap \gamma$. The rays $\{t\gamma \mid t \in \mathbb{R}^+, \gamma \in \mathcal{S}\}$ determined by simple closed curves are dense in $\mathcal{ML}(S)$.

The space $\mathcal{ML}(S)$ is a contractible topological manifold homeomorphic to $\mathbb{R}^{6g-6}$, but it does not have a natural smooth structure. Its natural structure is that of a piecewise linear (PL) manifold, with charts corresponding to train tracks.

Detailed discussion of the space $\mathcal{ML}(S)$ can be found in [Thu1] [EM] [PH] [Ota].

Grafting. As mentioned in the introduction, Thurston showed that grafting along simple closed curves has a natural extension to measured laminations, giving a projective grafting homeomorphism $\text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{P}(S)$ and a conformal grafting map $\text{gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{T}(S)$. Tanigawa showed that the latter is a proper map when either one of the two parameters is fixed:

Theorem 2.1 (Tanigawa [Tan]). *For any $\lambda \in \mathcal{ML}(S)$, the λ -grafting map $\text{gr}_\lambda : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is proper. For any $X \in \mathcal{T}(S)$, the X -grafting $\text{gr}_X : \mathcal{ML}(S) \rightarrow \mathcal{T}(S)$ is proper.*

The properness of the restricted grafting maps is used in the proofs of Theorems 1.1 (in §4) and 1.2 (in [SW]) to reduce a global statement to a local one, which is then attacked using infinitesimal methods. In the case of λ -grafting, the infinitesimal analysis is possible because $\text{gr}_\lambda : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is differentiable, and even real-analytic [McM, Cor. 2.11]. (A related real-analyticity property along rays in $\mathcal{ML}(S)$ is discussed in §6 below.)

The main step in [SW] is to show that the differential map $d\text{gr}_\lambda : T_X \mathcal{T}(S) \rightarrow T_{\text{gr}_\lambda X} \mathcal{T}(S)$ is an isomorphism. Once that is established, Theorem 1.2 follows easily, since gr_λ is then a proper local diffeomorphism of $\mathcal{T}(S)$, hence a covering map of the simply connected space $\mathcal{T}(S)$.

We will follow an analogous outline in the proof of Theorem 1.1, but the infinitesimal analysis is complicated by lack of smooth structure on $\mathcal{ML}(S)$, so the derivative of X -grafting does not exist in the classical sense. Instead we must use a weaker notion of differentiability based on one-sided derivatives, which we now discuss.

Tangentiability. A *tangentiability* map¹ $f : U \rightarrow V$ between open sets in $\mathbb{R}^n$ is a map with “one-sided” directional derivatives everywhere; in other words, for each $x \in U$ and $v \in \mathbb{R}^n$, the limit

$$(2.1) \quad \left. \frac{d}{dt} \right|_{t=0^+} f(x + tv) = \lim_{t \rightarrow 0^+} \frac{f(x + tv) - f(x)}{t}$$

exists, and the convergence is locally uniform in v (for equivalent conditions, see [Bon5, §2]). This convergence allows us to define $T_x f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, the *tangent map of f at x* , by

$$T_x f(v) = \left. \frac{d}{dt} \right|_{t=0^+} f(x + tv)$$

Of course if f is differentiable, then $T_x f$ is just the derivative of f at x , a linear map. When f is only tangentiability, the map $T_x f$ is continuous and homogeneous in the sense that $T_x f(\lambda v) = \lambda T_x f(v)$ for $\lambda \in \mathbb{R}^+$ [Bon5, §1].

A *tangentiability manifold* is one whose transition functions are tangentiability maps; examples include smooth manifolds and PL manifolds. Thus $\mathcal{ML}(S)$ has a natural tangentiability structure. The tangent space $T_x M$ at a point x of a tangentiability manifold is not naturally a vector space, but has the structure of a cone. The notion of tangentiability map extends naturally to tangentiability manifolds.

We will say that a homeomorphism between two tangentiability manifolds is a *bitangentiability homeomorphism* if it and its inverse are tangentiability, and if the tangent maps are everywhere homeomorphisms. A convenient criterion for this is provided by the

Lemma 2.2 (Bonahon [Bon5, Lem. 4]). *Let $f : M \rightarrow N$ be a homeomorphism between tangentiability manifolds. If f is tangentiability, and all of its tangent maps are injective, then f is a bitangentiability homeomorphism.*

Bonahon showed that grafting is compatible with that tangentiability structure of $\mathcal{ML}(S)$ in the following sense:

¹Some authors say instead that the map is *one-sided Gateaux differentiable*.

Theorem 2.3 (Bonahon [Bon5, Thm. 3]). *The grafting map $\text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{P}(S)$ is a bitangential homeomorphism. In particular, the conformal grafting map $\text{gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is tangential, and for each $X \in \mathcal{T}(S)$, the X -grafting map $\text{gr}_\bullet X : \mathcal{ML}(S) \rightarrow \mathcal{T}(S)$ is bitangential.*

In [Bon5], Bonahon actually computes the tangent map of grafting to show that grafting is tangential. After developing the shearing coordinates in §3, Bonahon's description of the tangent map (from which Theorem 2.3 is derived) is given in Theorem 3.4.

Differentiability. A curious feature of the tangentiality of grafting with respect to $\mathcal{ML}(S)$ is that some fragments of classical differentiability remain. For example, the inverse of the projective grafting map $\text{Gr}^{-1} : \mathcal{P}(S) \rightarrow \mathcal{ML}(S) \times \mathcal{T}(S)$ factors into the two projections $p_{\mathcal{ML}}$ and $p_{\mathcal{T}}$ (as described in the introduction). By Theorem 2.3, these are also tangential maps, but since both the domain and range of $p_{\mathcal{T}}$ are smooth manifolds, it makes sense to ask if this map is differentiable in the usual sense. Extending Theorem 2.3, Bonahon shows

Theorem 2.4 (Bonahon [Bon5, Thm. 3]). *The map $p_{\mathcal{T}} : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ is C^1 .*

In the same article, Bonahon shows that $p_{\mathcal{T}}$ fails to be C^2 for a certain family of punctured torus groups, which suggests that $p_{\mathcal{T}}$ may fail to be C^2 for all Teichmüller spaces $\mathcal{T}(S)$.

Finally we observe that $p_{\mathcal{T}}(\text{Gr}_\lambda X) = X$, so for each $\lambda \in \mathcal{ML}(S)$ we have $p_{\mathcal{T}} \circ \text{Gr}_\lambda = \text{Id}$, that is, the map $\text{Gr}_\lambda : \mathcal{T}(S) \rightarrow \mathcal{P}(S)$ is a smooth section of $p_{\mathcal{T}}$. Since $\text{Gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{P}(S)$ is a homeomorphism, these sections fill up $\mathcal{P}(S)$, and we conclude

Corollary 2.5. *The map $p_{\mathcal{T}}$ is a C^1 submersion.*

Infinitesimal X -grafting. Using the tangentiality of grafting, we can formulate the infinitesimal statement that will be our main tool in the proof of Theorem 1.1:

Theorem 2.6. *The tangent map $T_\lambda \text{gr}_\bullet X$ of the X -grafting map has no kernel. That is, if λ_t is a tangential family of measured laminations and $\frac{d}{dt} \big|_{t=0+} \text{gr}_{\lambda_t} X = 0$, then $\frac{d}{dt} \big|_{t=0+} \lambda_t = 0$.*

In the next two sections, we develop machinery to prove this result about the derivative of grafting, then strengthen it to a local injectivity result in order to prove Theorem 1.1. Complications arise in both steps because the maps under consideration are tangential rather than smooth.

3. SHEARING

In this section we describe the machinery of shearing cocycles for geodesic laminations on a hyperbolic surface, borrowing heavily from the papers of Bonahon [Bon4], [Bon2]. Some examples and discussion are included here to clarify the technicalities that our later arguments will require.

Cocycles. Let G be an abelian group and let $\Lambda \in \mathcal{GL}(S)$. A G -valued cocycle on Λ is a map α that assigns to each transversal τ to Λ an element $\alpha(\tau) \in G$ in a manner compatible with splitting and transversality-preserving isotopy. The G -module of all G -valued cocycles on Λ is denoted $\mathcal{H}(\Lambda, G)$.

Of particular interest for our purposes is the vector space of cocycles for maximal laminations with values in $\mathbb{R}$. While it is perhaps not clear from the definition, this vector space is finite dimensional, and the dimension is the same for all maximal laminations:

Theorem 3.1 (Bonahon [Bon2, Prop. 1]). *Let $\Lambda \in \mathcal{GL}(S)$ be a maximal lamination. Then $\mathcal{H}(\Lambda, \mathbb{R}) \simeq \mathbb{R}^{6g-6}$.*

The vector space $\mathcal{H}(\Lambda, \mathbb{R})$ carries a natural alternating bilinear form $\omega : \mathcal{H}(\Lambda, \mathbb{R}) \times \mathcal{H}(\Lambda, \mathbb{R}) \rightarrow \mathbb{R}$, the *Thurston symplectic form*, which comes from the cup product on $H^1(S, \mathbb{R})$ (see [PH] [Bon2, §3] [SB]). When Λ is maximal, the form ω is nondegenerate, making $\mathcal{H}(\Lambda, \mathbb{R})$ a symplectic vector space. While we do not use this symplectic structure directly in the proof of the main theorem, it is relevant to some of the constructions and examples in the sequel.

If $\lambda \in \mathcal{ML}(S)$ is a measured lamination with $\text{supp}(\lambda) \subset \Lambda \in \mathcal{GL}(S)$, then the total measure $\tau \mapsto \lambda(\tau)$ defines a real-valued cocycle on Λ , which we also denote by $\lambda \in \mathcal{H}(\Lambda, \mathbb{R})$. Cocycles arising from measures in this way take only nonnegative values on transversals; Bonahon showed that the converse is also true.

Theorem 3.2 (Bonahon [Bon4, Prop. 18]). *A transverse cocycle $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$ arises from a transverse measure for Λ if and only if $\alpha(\tau) \geq 0$ for every transversal τ .*

We therefore define $\mathcal{M}(\Lambda) \subset \mathcal{H}(\Lambda, \mathbb{R})$, the *cone of transverse measures* for Λ , to be the set of cocycles α satisfying $\alpha(\tau) \geq 0$ for all transversals τ . The set $\mathcal{M}(\Lambda)$ is a convex cone in the vector space $\mathcal{H}(\Lambda, \mathbb{R})$.

While positive real-valued cocycles on Λ correspond to transverse measures, there is an essential difference between a real-valued cocycle $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$ (whose value on a transversal is a real number) and a *signed transverse measure* on Λ which assigns a countably additive signed measure to each transversal τ (see Examples 1-2 below).

An analogous situation is the set function $[a, b] \mapsto (f(b) - f(a))$, where f is a real-valued function; this is a finitely additive function on intervals, but it only arises from a signed Borel measure if f has bounded variation (which is automatic if f is monotone). The connection between this example and real-valued cocycles for a lamination can be seen through the “distribution function” $f(x) = \alpha(\tau_x)$ where $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$, $\tau : [0, 1] \rightarrow S$ is a transversal, and $\tau_x = \tau|_{[0, x]}$. This function is defined for a.e. $x \in [0, 1]$ and monotonicity (for every τ) is equivalent to α being nonnegative.

The difference between measures and cocycles is also apparent from the dimension of the span of $\mathcal{M}(\Lambda)$, which is often 1 (by the solution to the Keane conjecture, see [Mas2] [Vee] [Ree] [Ker1]) and is never more than $(3g - 3)$ (because for maximal Λ , the space $\mathcal{H}(\Lambda, \mathbb{R}) \simeq \mathbb{R}^{6g-6}$ is a symplectic vector space in which $\text{span}(\mathcal{M}(\Lambda))$ is isotropic, see [Pap] [Lev] [Kat]). It follows that $\mathcal{M}(\Lambda)$ has positive codimension when, for example, Λ is maximal.

Shearing coordinates. Let $\Lambda \in \mathcal{GL}(S)$ be a maximal lamination, realized as a partial foliation of $X \in \mathcal{T}(S)$ by hyperbolic geodesics. Its lift $\tilde{\Lambda}$ to the universal cover $\tilde{X} \simeq \mathbb{H}^2$ determines a (not necessarily locally finite) tiling of $\mathbb{H}^2$ by ideal triangles.

A transversal $\tau : [a, b] \rightarrow \mathbb{H}^2$ to $\tilde{\Lambda}$ determines a pair of ideal triangles T_a and T_b which are the complementary regions of $\tilde{\Lambda}$ containing $\tau(a)$ and $\tau(b)$. In [Bon2], Bonahon constructs a *shearing cocycle* $\sigma(X) = \sigma^\Lambda(X) \in \mathcal{H}(\Lambda, \mathbb{R})$ from these data with the property that $\sigma(X)(\tau)$ measures the “relative shear” of the triangles T_a and T_b in $\mathbb{H}^2$. For example, the relative shear of two ideal triangles that share an edge is the signed distance between the feet of the altitudes based on the common side.

Remarkably, $\sigma(X)$ determines the metric X , and the set of such cocycles admits an explicit description. Let $\mathcal{C}(\Lambda) \subset \mathcal{H}(\Lambda, \mathbb{R})$ denote the set of $\mathbb{R}$ -valued cocycles that arise as shearing cocycles of hyperbolic metrics, and recall that $\omega : \mathcal{H}(\Lambda, \mathbb{R}) \times \mathcal{H}(\Lambda, \mathbb{R}) \rightarrow \mathbb{R}$ is the Thurston symplectic form.

Theorem 3.3 (Bonahon [Bon2, Thm. A,B]). *A cocycle $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$ is the shearing cocycle of a hyperbolic metric if and only if $\omega(\alpha, \mu) > 0$ for all $\mu \in \mathcal{M}(\Lambda)$, and $\mathcal{C}(\Lambda)$ is an open convex cone with finitely many faces. Furthermore $\sigma : \mathcal{T}(S) \rightarrow \mathcal{C}(\Lambda)$ is a real-analytic diffeomorphism.*

The condition $\omega(\alpha, \mu) > 0$ in Theorem 3.3 is necessary because for each $\mu \in \mathcal{M}(\Lambda)$, the Thurston pairing $\omega(\sigma(X), \mu)$ is the hyperbolic length of μ on X [Bon2, Thm. 9].

While the convex cone $\mathcal{C}(\Lambda)$ has finitely many faces and is a union of open rays in $\mathcal{H}(\Lambda, \mathbb{R})$, the zero cocycle $0 \in \mathcal{H}(\Lambda, \mathbb{R})$ is *not* an extreme point of $\mathcal{C}(\Lambda)$. In fact, the vector space of signed transverse measures (i.e. $\text{span } \mathcal{M}(\Lambda)$) is ω -isotropic (see [Pap]), and therefore

$$(3.1) \quad \text{span } \mathcal{M}(\Lambda) \subset \partial \mathcal{C}(\Lambda).$$

A schematic representation $\mathcal{C}(\Lambda)$ appears in Figure 2, where $\partial \mathcal{C}(\Lambda)$ contains a one-dimensional subspace of $\mathcal{H}(\Lambda, \mathbb{R})$.

Remark. Thurston parametrizes $\mathcal{T}(S)$ by a convex cone of shear coordinates in §9 of [Thu3]. The setting is in terms of duals to weights on train tracks, but the results are equivalent to those in Theorem 3.3.

Shearing maps. We now use the shearing embedding $\sigma : \mathcal{T}(S) \rightarrow \mathcal{H}(\Lambda, \mathbb{R})$ to turn translation in the vector space $\mathcal{H}(\Lambda, \mathbb{R})$ into a (locally-defined) map of Teichmüller space.




FIGURE 2. The shearing map is a translation in the shearing embedding of $\mathcal{T}(S)$ in $\mathcal{H}(\Lambda, \mathbb{R})$.

Let $X \in \mathcal{T}(S)$ and $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$. If the sum $\sigma(X) + \alpha$ is the shearing cocycle of a hyperbolic surface, we call this hyperbolic surface $\text{sh}_\alpha X$, the *shearing of X by α* . Thus $\text{sh}_\alpha X$ is defined by the condition

$$(3.2) \quad \sigma(\text{sh}_\alpha X) = \sigma(X) + \alpha.$$

Since $\sigma(\mathcal{T}(S)) = \mathcal{C}(\Lambda)$ is open, there is a neighborhood $U \subset \mathcal{H}(\Lambda, \mathbb{R}) \times \mathcal{T}(S)$ of $\{0\} \times \mathcal{T}(S)$ in which the shearing map $\text{sh} : U \rightarrow \mathcal{T}(S)$ is well-defined.

In particular, for any fixed $X \in \mathcal{T}(S)$ and $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$ there is some $\epsilon > 0$ such that $\text{sh}_{t\alpha} X$ is defined for all $|t| < \epsilon$ (see Figure 2), and

$$\frac{d}{dt} \sigma(\text{sh}_{t\alpha} X) = \alpha.$$

We consider a few examples of shearing maps to highlight the role of $\mathcal{C}(\Lambda)$ and the fact that $\text{sh}_\alpha X$ is not defined for all pairs (α, X) . First, if Λ contains a simple closed geodesic γ , then for each $t \in \mathbb{R}$ the cocycle $t\gamma$ is a signed transverse measure. Furthermore, for all $X \in \mathcal{T}(S)$, we have $(\sigma(X) + t\gamma) \in \mathcal{C}(\Lambda)$, since $t\gamma \in \mathcal{M}(\Lambda)$ and the span of $\mathcal{M}(\Lambda)$ is ω -isotropic; thus $\text{sh}_{t\gamma} X$ is defined for all $t \in \mathbb{R}$. Concretely, the hyperbolic surface $\text{sh}_{t\gamma} X$ is obtained from X by cutting along the geodesic γ and then gluing the two boundary components with a twist (by signed distance t).

This twisting example has a natural generalization: given a cocycle $\lambda \in \mathcal{H}(\Lambda, \mathbb{R})$ representing a transverse measure for Λ , the shearing $\text{sh}_{t\lambda}(X)$ is again defined for all $t \in \mathbb{R}$ and the resulting map $\text{sh}_{t\lambda} : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is called an *earthquake*. For further discussion of earthquakes, see [Thu2] [Ker3] [EM] [Bon1] [McM].

As a final example, consider shearing a surface X using a cocycle $\alpha \in \mathcal{H}(\Lambda, \mathbb{R})$ that is itself the shearing cocycle of a hyperbolic surface, i.e. $\alpha \in \mathcal{C}(\Lambda)$. For any transverse measure $\lambda \in \mathcal{M}(\Lambda)$, the hyperbolic length of λ on $\text{sh}_{t\alpha} X$ can be computed using the Thurston intersection form ω (cf. [Bon2, Thm. 9]), and we have

$$(3.3) \quad \ell(\lambda, \text{sh}_{t\alpha} X) = \omega(\sigma(X) + t\alpha, \lambda) = \ell(\lambda, X) + t\omega(\alpha, \lambda) = A + Bt$$

where $A, B > 0$. Since the length of a measured lamination is positive, it follows that the set of t for which $\sigma(X) + t\alpha$ is the shearing cocycle of a hyperbolic metric (and hence those for which $\text{sh}_{t\alpha} X$ exists) is a subset of $\{t > -(A/B)\}$.

Tangent cocycles. Let $\lambda_t \in \mathcal{ML}(S)$, $t \in [0, \epsilon]$ be a tangentiable ray of measured laminations. We will represent the tangent vector $\frac{d}{dt}|_{t=0^+} \lambda_t$ by a transverse cocycle to a certain geodesic lamination (as in [Bon3]). We first describe the underlying geodesic lamination.

The *essential support* of λ_t at $t = 0^+$ is a geodesic lamination $\Lambda \in \mathcal{GL}(S)$ that reflects how the support of λ_t is changing for small positive values of t . For a *PL* family of measured laminations λ_t , the essential support is the Hausdorff limit $\lim_{t \rightarrow 0^+} \text{supp}(\lambda_t)$ of the supporting geodesic laminations [Bon3, Prop. 4]. For the general case, we only sketch the construction, and refer the reader to [Bon3, §2] for details.

First lift λ_t to a family $\tilde{\lambda}_t$ of measured geodesic laminations in $\mathbb{H}^2$. Define a set $\tilde{\Lambda}$ of geodesics in $\mathbb{H}^2$ as follows: a geodesic γ belongs to $\tilde{\Lambda}$ if and only if for every smooth transversal $\tau : [-\epsilon, \epsilon] \rightarrow \mathbb{H}^2$ with $\tau(0) \in \gamma$, the total transverse measure of τ with respect to λ_t is at least Ct for some $C > 0$ (depending on τ and γ) and all t sufficiently small. Then $\tilde{\Lambda}$ is the lift of a geodesic lamination $\Lambda \in \mathcal{GL}(S)$, the essential support of λ_t .

The tangent vector $\frac{d}{dt}|_{t=0^+} \lambda_t$ defines a real-valued transverse cocycle $\dot{\lambda} \in \mathcal{H}(\Lambda, \mathbb{R})$ on the essential support as follows:

$$\dot{\lambda}(\tau) = \lim_{t \rightarrow 0^+} \frac{1}{t} (\lambda_t(\tau) - \lambda_0(\tau)).$$

This is the *tangent cocycle* of λ_t at $t = 0^+$. Clearly the same formula defines a cocycle for a lamination containing the essential support of λ_t , so as a convenience we may assume Λ is maximal. We illustrate this construction with two examples:

Example 1. The cocycle determined by a measured lamination $\lambda \in \mathcal{M}(\Lambda)$ is the tangent cocycle of the “ray” $\lambda_t = (1 + t)\lambda \in \mathcal{ML}(S)$ at $t = 0^+$.

Example 2. Let S be a punctured torus with meridian α and longitude β (putting aside our assumption that S is compact for a moment), and consider the family $\lambda_t \in \mathcal{ML}(S)$ defined by the conditions $\lambda_t(\alpha) = 1$ and $\lambda_t(\beta) = t$. Thus for each $n \in \mathbb{N}$, the measured lamination $\lambda_{1/n}$ is a simple closed curve with homology class $(n[\alpha] + [\beta])$ and weight $1/n$.

The essential support of λ_t at $t = 0^+$ is the geodesic lamination $\Lambda = \alpha \cup \eta$, where η is an infinite simple geodesic that spirals toward α in each direction. The derivative $\dot{\lambda} \in \mathcal{H}(\Lambda, \mathbb{R})$ is a cocycle with full support and indefinite sign (compare Bonahon's example [Bon3, p. 104]).

This second example shows that the tangent cocycle may not be a transverse measure, and the essential support may not admit a measure of full support. This illustrates some of the difficulties of using differential methods on the space of measured laminations and of adapting the methods in [SW] to prove the main theorem.

Following Thurston, Bonahon showed that the association of a cocycle $\dot{\lambda}$ to a tangentiable family λ_t provides a linear model for each PL face of the tangent space $T_{\lambda_0} \mathcal{ML}(S)$ (see [Bon3], [Thu3, §6]). When $\text{supp}(\lambda_0)$ is not maximal, however, no single maximal lamination Λ can be chosen to contain the essential support of *every* family λ_t , and so there is no embedding $T_{\lambda_0} \mathcal{ML}(S) \rightarrow \mathcal{H}(\Lambda, \mathbb{R})$.

In contrast, when $\text{supp}(\lambda_0) = \Lambda$ is maximal (a generic situation that excludes, for example, closed leaves), the tangent cocycle construction defines a homeomorphism $T_{\lambda_0} \mathcal{ML}(S) \simeq \mathcal{H}(\Lambda, \mathbb{R})$, giving the tangent space a canonical linear structure. This was observed in [Thu3].

Complex linearity. So far we have seen real-valued cocycles on geodesic laminations arise in two different contexts: first as shearing cocycles providing coordinates for $\mathcal{T}(S)$, and then as tangent vectors to families of measured laminations. Building on these two constructions, the following result of Bonahon will allow us to connect the derivative of grafting with respect to $\mathcal{T}(S)$ and $\mathcal{ML}(S)$:

Theorem 3.4 (Bonahon [Bon5, Prop. 5] [Bon2, §10]). *Let $X \in \mathcal{T}(S)$ and $\lambda \in \mathcal{ML}(S)$. For each maximal geodesic lamination $\Lambda \in \mathcal{GL}(S)$ containing the support of λ , there is a complex-linear map $L = L(\Lambda, \lambda, X) : \mathcal{H}(\Lambda, \mathbb{C}) \rightarrow T_{\text{Gr}_\lambda X} \mathcal{P}(S)$ which determines the tangent map of Gr in tangent directions carried by Λ , in the following sense:*

Let $\lambda_t \in \mathcal{ML}(S)$ be a tangentiable family of measured laminations with $\lambda_0 = \lambda$ and with essential support contained in Λ , and let $\dot{\lambda} = \frac{d}{dt}|_{t=0^+} \lambda_t \in \mathcal{H}(\Lambda, \mathbb{R})$. Let $X_t \in \mathcal{T}(S)$ be a smooth family of hyperbolic structures with $X_0 = X$ whose derivative in the Λ shearing embedding is $\dot{\sigma} = \frac{d}{dt}|_{t=0^+} \sigma(X_t)$.

Then $t \mapsto \text{Gr}_{\lambda_t} X_t$ is a tangentiable curve in $\mathcal{P}(S)$ and

$$\left. \frac{d}{dt} \right|_{t=0^+} \text{Gr}_{\lambda_t} = L(\dot{\sigma} + i\dot{\lambda})$$

Similarly $\text{gr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is tangentiable and its derivative has the same complex linearity property.

In terms of the piecewise linear structure of $\mathcal{ML}(S)$, Theorem 3.4 says that on each linear face of the tangent space $T_{(\lambda, X)}(\mathcal{ML}(S) \times \mathcal{T}(S))$, the

tangent map $T_{(\lambda, X)} \text{gr}$ is the restriction of a complex-linear map $\mathcal{H}(\Lambda, \mathbb{C}) \rightarrow T_{\text{gr}_\lambda X} \mathcal{T}(S)$ (see [Bon5, §2] [Bon2, §10]).

4. PROOF OF THE MAIN THEOREM

With the necessary background in place, we can now show that the tangent map of $\text{gr}_\bullet X$ has no kernel.

Proof of Theorem 2.6. Fix X and suppose that $\frac{d}{dt}|_{t=0^+} \text{gr}_{\lambda_t} X = 0$. Let $\dot{\lambda} \in \mathcal{H}(\Lambda, \mathbb{R})$ denote the derivative of λ_t at $t = 0^+$, with $\Lambda \in \mathcal{GL}(S)$ maximal. We must show that $\dot{\lambda} = 0$.

For all t sufficiently small, the shearing $X_t = \text{sh}_{t\dot{\lambda}} X \in \mathcal{T}(S)$ is defined and satisfies $X_0 = X$ and $\dot{\lambda} = \frac{d}{dt} \sigma(X_t)$. Therefore by Theorem 3.4,

$$i \left[\frac{d}{dt} \Big|_{t=0^+} \text{gr}_{\lambda_0} X_t \right] = iL(\dot{\lambda}) = L(i\dot{\lambda}) = \frac{d}{dt} \Big|_{t=0^+} \text{gr}_{\lambda_t} X_0 = 0,$$

where L is the complex-linear map representing the tangent map of gr in tangent directions carried by Λ .

By Theorem 1.2, gr_{λ_0} is an immersion, and so $\frac{d}{dt}|_{t=0^+} X_t = 0$. Thus

$$\dot{\lambda} = \frac{d}{dt} \sigma(X_t) = 0.$$

□

If the grafting map were continuously differentiable in the usual sense, the proof of the main theorem would now be straightforward, using linearity of the derivative and the inverse function theorem to conclude that $\text{gr}_\bullet X$ is a local diffeomorphism. We will follow this general outline, but we will need to use additional properties of the grafting maps to strengthen the infinitesimal result to a local one.

In fact, some argument specific to grafting is necessary at this point. In general, a tangent map that has no kernel need not be injective (consider $\mathbb{C} \rightarrow \mathbb{R}^+$ by $z \mapsto |z|$ at $z = 0$), and even if the tangent map is injective, tangentiability does not imply the continuous variation of derivatives needed for the inverse function theorem.

With these potential problems in mind, we analyze the X -grafting map $\text{gr}_\bullet X$ as the composition of the projective X -grafting map $\text{Gr}_\bullet X$ and the smooth projection $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$. Let $N = \dim \mathcal{T}(S) = \dim \mathcal{ML}(S) = \frac{1}{2} \dim \mathcal{P}(S) = 6g - 6$.

Let $G_X = \{\text{Gr}_\lambda X \mid \lambda \in \mathcal{ML}(S)\} \subset \mathcal{P}(S)$ denote the image of $\text{Gr}_\bullet X$. Recall the map $p_{\mathcal{T}} : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ is defined by $p_{\mathcal{T}}(\text{Gr}_\lambda X) = X$, and so $G_X = p_{\mathcal{T}}^{-1}(X)$ is a fiber of this map. Since $p_{\mathcal{T}}$ is a C^1 submersion (by Corollary 2.5), the set G_X is actually a C^1 submanifold of $\mathcal{P}(S)$ of dimension $N = \dim \mathcal{P}(S) - \dim \mathcal{T}(S)$. In particular, G_X is smoother than its tangentiability parameterization by $\mathcal{ML}(S)$ would suggest.

We can recast Theorem 2.6 as a result about the tangent space to G_X as follows:

Theorem 4.1. *For any $X \in \mathcal{T}(S)$, the C^1 submanifold $G_X \subset \mathcal{P}(S)$ is transverse to the map $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$; that is, for any $Z \in G_X$, we have $T_Z G_X \cap (\ker d\pi) = \{0\}$. Equivalently, the distributions $\ker d\pi$ and $\ker dp_{\mathcal{T}}$ in $T\mathcal{P}(S)$ are transverse.*

Proof. Suppose not, i.e. that there exists $v \in (T_Z G_X \cap \ker d\pi)$ with $v \neq 0$. Since $Z \in G_X$, we have $Z = \text{Gr}_{\lambda} X$ for some $\lambda \in \mathcal{ML}(S)$. Choose a C^1 path Z_t in G_X with $Z_0 = Z$ and $\frac{d}{dt}|_{t=0+} Z_t = v$, and let $\text{Gr}^{-1}(Z_t) = (\lambda_t, X)$. Since Gr^{-1} is a bitangentiable homeomorphism, the family λ_t is tangentiable and satisfies $\frac{d}{dt}|_{t=0+} \lambda_t \neq 0$. On the other hand, $\frac{d}{dt}|_{t=0+} \text{gr}_{\lambda_t} X = \frac{d}{dt}|_{t=0+} \pi(Z_t) = d\pi(\frac{d}{dt}|_{t=0+} Z_t) = 0$, contradicting Theorem 2.6. $\square$

While a general tangent map can have no kernel and yet fail to be injective, the fact that G_X is C^1 (and thus has linear tangent spaces) rules out this behavior for $\text{Gr}_{\bullet} X$:

Theorem 4.2. *The tangent map $T_{\lambda} \text{gr}_{\bullet} X$ of the conformal X -grafting map is a homeomorphism.*

Proof. We study the diagram of grafting maps:

$$\begin{array}{ccc} \mathcal{ML}(S) & \xrightarrow{\text{Gr}_{\bullet} X} & G_X \subset \mathcal{P}(S) \\ & \searrow \text{gr}_{\bullet} X & \downarrow \pi \\ & & \mathcal{T}(S) \end{array}$$

Fix $\lambda \in \mathcal{ML}(S)$ and for brevity let $f = T_{\lambda} \text{gr}_{\bullet} X$ be the tangent map of the conformal X -grafting map; our goal is to show that f is a homeomorphism.

Similarly, let $F = T_{\lambda} \text{Gr}_{\bullet} X$ be the tangent map of the projective X -grafting map. Then $f = d\pi \circ F$, where $d\pi$ is the differential of $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ at $Z = \text{Gr}_{\lambda} X$, i.e. the tangent maps form a corresponding diagram:

$$\begin{array}{ccc} T_{\lambda} \mathcal{ML}(S) & \xrightarrow{F} & T_Z G_X \subset T_Z \mathcal{P}(S) \\ & \searrow f & \downarrow d\pi \\ & & T_{\pi(Z)} \mathcal{T}(S) \end{array}$$

Note that F and f are continuous homogeneous maps, while $d\pi$ is linear.

Since Gr is a tangentiable homeomorphism, its restriction $\text{Gr}_{\bullet} X : \mathcal{ML}(S) \rightarrow \mathcal{P}(S)$ is a tangentiable injection, i.e. its tangent map F is both injective and a homogeneous homeomorphism onto its image. Since G_X is a C^1 submanifold of $\mathcal{P}(S)$, the image of F is the linear subspace $T_Z G_X \subset T_Z \mathcal{P}(S)$.

First we show that f is injective. Suppose on the contrary there exist $\dot{\lambda}_1, \dot{\lambda}_2 \in T_{\lambda} \mathcal{ML}(S)$, distinct and nonzero, and $f(\dot{\lambda}_1) = f(\dot{\lambda}_2)$. Then $v = F(\dot{\lambda}_1) - F(\dot{\lambda}_2) \in T_Z G_X$ is nonzero since F is injective, and so $v \in \ker d\pi$, which contradicts Theorem 4.1.

Thus $f : T_\lambda \mathcal{ML}(S) \rightarrow T_{\text{gr}_\lambda X} \mathcal{T}(S)$ is injective. Since f is also a homogeneous map between cones of the same dimension, it is a homeomorphism. $\square$

Remark. The fact that $T_\lambda \text{Gr}_\bullet X$ is a homeomorphism onto a linear subspace of $T_{\text{Gr}_\lambda X} \mathcal{P}(S)$ is also observed by Bonahon in the proof of Prop. 12 in [Bon5].

Finally we complete our study of $\text{Gr}_\bullet X$ and $\text{gr}_\bullet X$ by proving the main theorem:

Proof of Theorem 1.1. Let us consider the restriction of the forgetful map $\pi : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ to the C^1 submanifold $G_X \subset \mathcal{P}(S)$. Since G_X is N -dimensional, this projection is a local diffeomorphism at $Z \in G_X$ if the subspaces $T_Z G_X$ and $\ker d\pi$ of $\mathcal{T}_Z \mathcal{P}(S)$ are transverse (by the inverse function theorem). By Theorem 4.1, this is true for every $Z \in G_X$, so $\pi|_{G_X}$ is a local C^1 diffeomorphism.

Thus the conformal X -grafting map is the composition of the homeomorphism $\text{Gr}_\bullet X : \mathcal{ML}(S) \rightarrow G_X$ and the local homeomorphism $\pi|_{G_X} : G_X \rightarrow \mathcal{T}(S)$, so $\text{gr}_\bullet X$ is a local homeomorphism. As we noted in Section 2, by the properness of $\text{gr}_\bullet X$ (Theorem 2.1), it follows that this map is a homeomorphism. Thus X -grafting is a tangential homeomorphism with injective tangent maps (Theorem 4.2), which by Lemma 2.2 is a bitangential homeomorphism. $\square$

5. APPLICATIONS

In this section, we collect some applications of the main theorem itself and of the techniques used in its proof, and we discuss some related questions about grafting coordinates and $\mathbb{CP}^1$ structures.

Projections. We begin by proving the main corollary of Theorems 1.1 and 1.2 about the grafting coordinates for a fiber $P(X) = \pi^{-1}(X) \subset \mathcal{P}(S)$:

Corollary 1.3. *For each $X \in \mathcal{T}(S)$, the space $P(X)$ of $\mathbb{CP}^1$ structures on X is a graph over each factor in the grafting coordinate system. In fact, we have:*

- (1) *The projection $p_{\mathcal{T}}|_{P(X)} : P(X) \rightarrow \mathcal{T}(S)$ is a C^1 diffeomorphism.*
- (2) *The projection $p_{\mathcal{ML}}|_{P(X)} : P(X) \rightarrow \mathcal{ML}(S)$ is a bitangential homeomorphism.*

Proof. First we consider the regularity of the maps. Bonahon showed that $p_{\mathcal{T}}$ is C^1 (Theorem 2.4), and $p_{\mathcal{ML}}$ is tangential because it is the composition of Gr^{-1} , a bitangential homeomorphism (Theorem 2.3), and the projection to one factor of a product of tangential manifolds.

- (1) *Proof that $p_{\mathcal{T}}|_{P(X)}$ is a diffeomorphism.* First of all, the map $p_{\mathcal{T}}|_{P(X)}$ is a homeomorphism because it has inverse map

$$Y \mapsto \text{Gr}_{(\text{gr}_\bullet X)^{-1}(Y)} Y$$

where the map $(\text{gr}_\bullet X)^{-1} : \mathcal{T}(S) \rightarrow \mathcal{ML}(S)$ exists by Theorem 1.1.

Thus it suffices to show that $p_{\mathcal{T}}|_{P(X)}$ is a local diffeomorphism. But the kernel of $dp_{\mathcal{T}}|_{P(X)}$ is the intersection of $\ker d\pi$ and $\ker dp_{\mathcal{T}}$ in $T\mathcal{P}(S)$, which is zero by Theorem 4.1. So the derivative of $p_{\mathcal{T}}|_{P(X)}$ is an isomorphism, and by the inverse function theorem this map is a local diffeomorphism.

- (2) *Proof that $p_{\mathcal{ML}}|_{P(X)}$ is a bitangentiable homeomorphism.* As in (1), we first show that $p_{\mathcal{ML}}|_{P(X)}$ is a homeomorphism by exhibiting an inverse map,

$$\lambda \mapsto \text{Gr}_{\lambda}(\text{gr}_{\lambda}^{-1}(Y))$$

where gr_{λ}^{-1} exists by Theorem 1.2.

By Lemma 2.2, we need only show that the tangent map of $p_{\mathcal{ML}}|_{P(X)}$ is everywhere injective. That this tangent map has no kernel also follows easily from Theorem 1.2, but to show injectivity we will use an argument modeled on the proofs of Theorems 2.6 and 4.2.

Suppose on the contrary that two distinct, nonzero tangent vectors $v_1, v_2 \in T_Z P(X)$ have the same image in $T_{\lambda} \mathcal{ML}(S)$ under the tangent map of $p_{\mathcal{ML}}$, where $Z = \text{Gr}_{\lambda} Y$. Then differentiable paths in $P(X)$ with tangent vectors v_1 and v_2 are mapped by Gr^{-1} to tangentiable paths $\text{Gr}_{\lambda_t^{(1)}} Y_t^{(1)}$ and $\text{Gr}_{\lambda_t^{(2)}} Y_t^{(2)}$, respectively, where $Y_0^{(k)} = Y$ and $\lambda_0^{(k)} = \lambda$ for $k = 1, 2$.

In this notation, the image of v_k by the tangent map $T_Z p_{\mathcal{ML}}$ is $\frac{d}{dt}|_{t=0+} \lambda_t^{(k)}$, so we have $\frac{d}{dt}|_{t=0+} \lambda_t^{(1)} = \frac{d}{dt}|_{t=0+} \lambda_t^{(2)} = \dot{\lambda}$. In particular there is a single geodesic lamination Λ containing the essential support of both families $\lambda_t^{(k)}$, and their common tangent vector defines a cocycle $\dot{\lambda} \in \mathcal{H}(\Lambda, \mathbb{R})$. Using the shearing embedding of Teichmüller space in $\mathcal{H}(\Lambda, \mathbb{R})$ gives cocycles $\dot{\sigma}_k = \frac{d}{dt}|_{t=0+} (\sigma^{\Lambda}(Y_t^{(k)}))$, and since $v_1 \neq v_2$ we have $\dot{\sigma}_1 \neq \dot{\sigma}_2$.

By Theorem 3.4, there is a complex-linear map $L : \mathcal{H}(\Lambda, \mathbb{C}) \rightarrow T_X \mathcal{T}(S)$ that gives the tangent map of Gr for tangent vectors to $\mathcal{ML}(S) \times \mathcal{T}(S)$ at (λ, X) representable by complex-valued cocycles on Λ , so $v_k = L(\dot{\sigma}_k + i\dot{\lambda})$. Now consider $\dot{\sigma} = \dot{\sigma}_1 - \dot{\sigma}_2 \neq 0$, which is the tangent vector to the shearing family $Y_t = \text{sh}_{t\dot{\sigma}} Y$; we have

$$\begin{aligned} \frac{d}{dt}\Big|_{t=0+} \text{Gr}_{\lambda} Y_t &= L(\dot{\sigma}) \\ &= L\left((\dot{\sigma}_1 + i\dot{\lambda}) - (\dot{\sigma}_2 + i\dot{\lambda})\right) \\ &= v_1 - v_2 \in T_Z P(X), \end{aligned}$$

and so $\frac{d}{dt}|_{t=0+} \text{gr}_{\lambda} Y_t = 0$, which by Theorem 1.2 implies that $\frac{d}{dt}|_{t=0+} Y_t = 0$, and so $\dot{\sigma} = \frac{d}{dt}|_{t=0+} \sigma^{\Lambda}(Y_t) = 0$, a contradiction. Thus the tangent map of $p_{\mathcal{ML}}$ is injective, as required.

□

Remark. Since Bonahon constructs an explicit example to show that the full projection map $p_{\mathcal{T}} : \mathcal{P}(S) \rightarrow \mathcal{T}(S)$ is not necessarily C^2 , it would be interesting to know if a similar construction could be used to show that $p_{\mathcal{T}}|_{P(X)}$ need not be C^2 .

Pruning. Since the λ -grafting map $\text{gr}_{\lambda} : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ is a homeomorphism (Theorem 1.2), there is an inverse map $\text{pr}_{\lambda} : \mathcal{T}(S) \rightarrow \mathcal{T}(S)$ which we call *pruning by λ* . Roughly speaking, grafting by λ inserts a Euclidean subsurface along the leaves of λ , and pruning by λ removes it. Allowing λ to vary, we obtain the pruning map $\text{pr} : \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathcal{T}(S)$, and fixing X we have the X -pruning map $\text{pr}_{\cdot} X : \mathcal{ML}(S) \times \{X\} \rightarrow \mathcal{T}(S)$.

We can reformulate Corollary 1.3 in terms of pruning as follows:

Corollary 5.1. *For each $X \in \mathcal{T}(S)$, the X -pruning map $\text{pr}_{\cdot} X : \mathcal{ML}(S) \rightarrow \mathcal{T}(S)$ is a bitangentiable homeomorphism.*

Proof. The graph of the X -pruning map consists of the pairs (λ, Y) such that $\text{gr}_{\lambda} Y = X$, which is simply the fiber $P(X) \subset \mathcal{P}(S) \simeq \mathcal{ML}(S) \times \mathcal{T}(S)$. In terms of the projections $p_{\mathcal{ML}} : P(X) \rightarrow \mathcal{ML}(S)$ and $p_{\mathcal{T}} : P(X) \rightarrow \mathcal{T}(S)$, we have $\text{pr}_{\cdot} X = p_{\mathcal{T}} \circ p_{\mathcal{ML}}^{-1}$, which is a bitangentiable homeomorphism by Corollary 1.3. $\square$

Previously it was known that $\text{pr}_{\cdot} X$ is a “rough homeomorphism”, i.e. a proper map of degree 1, and that it has a natural extension to the Thurston compactification of $\mathcal{T}(S)$ and the projective compactification of $\mathcal{ML}(S)$ by $\mathbb{PML}(S) = (\mathcal{ML}(S) - \{0\})/\mathbb{R}^+$. Furthermore, the resulting boundary map $\mathbb{PML}(S) \rightarrow \mathbb{PML}(S)$ is the *antipodal map* relative to X . For details, see [D1].

Internal Coordinates for the Bers slice. Let $\mathcal{QF} = \mathcal{QF}(S)$ denote the space of marked quasi-Fuchsian hyperbolic 3-manifolds homeomorphic to $S \times \mathbb{R}$ (see [Ber1] [Nag]). Each such manifold $M \in \mathcal{QF}$ has ideal boundary $\partial_{\infty} M = Y_+ \sqcup Y_-$ with conformal structures $Y_{\pm} \in \mathcal{T}(S)$ and convex core boundary surfaces $X_{\pm} \in \mathcal{T}(S)$ with bending laminations $\lambda_{\pm} \in \mathcal{ML}(S)$ (represented schematically in Figure 3). Furthermore, the convex core and ideal boundary surfaces satisfy $\text{gr}_{\lambda_{\pm}} X_{\pm} = Y_{\pm}$ ([McM, Thm. 2.8], see also [KT]).

In his celebrated holomorphic embedding [Ber2] of the Teichmüller space $\mathcal{T}(S)$ into $\mathbb{C}^{3g-3}$, Bers focused on the slice $\mathcal{B}_Y \subset \mathcal{QF}$ of those quasi-Fuchsian manifolds with a fixed ideal boundary surface $Y_- = Y$. Bers showed that a quasi-Fuchsian manifold $M \in \mathcal{B}_Y$ is uniquely determined by its other conformal boundary surface $Y_+ \in \mathcal{T}(S)$, which can be chosen arbitrarily (the *Simultaneous Uniformization Theorem* [Ber1]).

A corollary of our main theorem is that $\mathcal{B}_Y$ may also be parameterized by the hyperbolic structure X_- :

Corollary 5.2. *Let $M, M' \in \mathcal{QF}$ be marked quasi-Fuchsian manifolds, and suppose an end of M (respectively M') has ideal boundary Y (resp. Y') and the associated convex core boundary surface has hyperbolic metric X (resp.*




FIGURE 3. Geometric data associated to a quasi-Fuchsian manifold $M \in \mathcal{QF}$. In a Bers slice, Y_- is fixed.

X'). If Y and Y' are conformally equivalent and X and X' are isometric, then M is isometric to M' .

Proof. By hypothesis $Y' = Y$ and $X' = X$ as points in Teichmüller space, so the bending measures λ and λ' of the convex core boundaries satisfy $\text{gr}_\lambda X = \text{gr}_{\lambda'} X = Y$. By Theorem 1.1, we have $\lambda = \lambda'$.

A quasi-Fuchsian manifold M is uniquely determined up to isometry by the hyperbolic metric X and bending lamination λ of one of its convex core boundary surfaces, since one can use X and λ to construct the associated equivariant pleated plane in $\mathbb{H}^3$ and its holonomy group $\pi_1 M \subset \text{PSL}_2(\mathbb{C})$ (see [EM]). As M and M' share these data, they are isometric. $\square$

For another perspective on this corollary, we can consider the Bers slice $\mathcal{B}_Y$ as a subset of $P(Y)$, where a quasi-Fuchsian manifold $M \in \mathcal{B}_Y$ is identified with the projective structure it induces on its ideal boundary surface Y . Then Corollary 5.2 is equivalent to the statement that for every $Y \in \mathcal{T}(S)$, the projection map $p_{\mathcal{T}}|_{\mathcal{B}_Y} : \mathcal{B}_Y \rightarrow \mathcal{T}(S)$ is injective. Of course this can also be derived from Corollary 1.3, which shows, furthermore, that $p_{\mathcal{T}}|_{\mathcal{B}_Y}$ is a C^1 embedding (since $\mathcal{B}_Y \subset P(Y)$ is open).

Remarks.

1. In [SW], it was observed that a manifold $M \in \mathcal{B}_Y$ is also determined by the bending lamination $\lambda_- \in \mathcal{ML}(S)$ on the same side as the fixed conformal structure Y . It is not known whether either the bending lamination λ_+ or the hyperbolic structure X_+ determine elements of $\mathcal{B}_Y$.
2. More generally, one can ask what geometric data determine $M \in \mathcal{QF}$ up to isometry. Bonahon and Otal [BO] showed that if λ_+ and λ_- bind the surface and are supported on simple closed curves, then the pair (λ_+, λ_-) determines $M \in \mathcal{QF}$ uniquely. Recently, Bonahon

[Bon6] showed that this restriction to simple closed curves may be lifted for elements of $\mathcal{QF}$ which are sufficiently close to the Fuchsian subspace $\mathcal{F} \subset \mathcal{QF}$, and Series [Ser3] proved that λ_+ and λ_- determine M when S is a once-punctured torus.

6. GRAFTING COORDINATES AND RAYS

In this final section we discuss how the ray structure of $\mathcal{ML}(S)$ is transported to $\mathcal{T}(S)$ by the X -grafting map.

Recall (from Section 2) that the action of $\mathbb{R}^+$ on $\mathcal{ML}(S)$ by scaling transverse measures gives this space the structure of a cone, with the empty lamination 0 as its base point. By Theorem 1.1, for each $X \in \mathcal{T}(S)$ we can use the X -grafting map to parameterize the Teichmüller space by $\mathcal{ML}(S)$, providing a global system of “polar coordinates” centered at X . In this coordinate system, the ray $\mathbb{R}^+\lambda \in \mathcal{ML}(S)$ corresponds to the *grafting ray* $\{\text{gr}_{t\lambda} X \mid t \in \mathbb{R}^+\}$, a properly embedded path starting at X , and Teichmüller space is the union of these rays.

It would be interesting to understand the geometry of this coordinate system, and especially the grafting rays. Thus we ask:

Question. What is the behavior of the grafting ray $t \mapsto \text{gr}_{t\gamma} X$ and how does it depend on λ and X ?

Theorems 6.2 and 6.6 below address this question by estimating the extremal and hyperbolic lengths of the grafting lamination (and their derivatives) along a ray. The asymptotic behavior of certain grafting rays in relation to the Teichmüller metric has also been investigated by Díaz and Kim, see [DK].

Naturally, one first wonders about the regularity of grafting rays, since the grafting map itself exhibits a combination of tangential and differentiable behavior. However, along rays the grafting map is as smooth as possible:

Theorem 6.1 (McMullen [McM]). *The grafting ray $t \mapsto \text{gr}_{t\lambda}$ is a real-analytic map from $\mathbb{R}^+$ to $\mathcal{T}(S)$; in fact, it is the restriction of a real-analytic map $\{t \geq -\epsilon\} \rightarrow \mathcal{T}(S)$ for some $\epsilon > 0$ (depending on λ). Furthermore, if $\lambda_n \rightarrow \lambda$ in $\mathcal{ML}(S)$, then the λ_n grafting rays converge in C^ω to the λ grafting ray.*

Remarks.

1. In [McM] it is shown that for any $\lambda \in \mathcal{ML}(S)$, the *complex earthquake map* $\text{eq}_\lambda : \mathbb{H} \rightarrow \mathcal{T}(S)$ is holomorphic and extends to an open neighborhood of $\mathbb{H} \cup \mathbb{R}$ (Proposition 2.6 and Theorem 2.10). Furthermore these maps vary continuously with $\lambda \in \mathcal{ML}(S)$ (Theorem 2.5), as do their derivatives, since they are holomorphic. Since the grafting ray is the restriction of the complex earthquake to $i\mathbb{R}$, Theorem 6.1 follows immediately.

2. The regularity of grafting rays and complex earthquakes is closely related to (and in part, an application of) the analyticity of quake-bend deformations of surface group representations in $\mathrm{PSL}_2(\mathbb{C})$ established by Epstein and Marden [EM].

Extremal length. When λ is supported on a simple closed geodesic γ , it is clear that a large grafting will result in a surface in which γ has small extremal length, because $\mathrm{gr}_{t\gamma} X$ contains an annulus of large modulus homotopic to γ . Refining this intuition, we will establish the estimate:

Theorem 6.2. *For each $X \in \mathcal{T}(S)$, the extremal length of λ is of order $1/t$ on the λ -grafting ray $t \mapsto \mathrm{gr}_{t\lambda} X$ and is monotone decreasing for all $t \gg 0$. Specifically, we have*

$$(1) \ E(\lambda, \mathrm{gr}_{t\lambda} X) = \frac{\ell(\lambda, X)}{t} + O(t^{-2}), \text{ and}$$

$$(2) \ \frac{d}{dt} E(\lambda, \mathrm{gr}_{t\lambda} X) = \frac{-\ell(\lambda, X)}{t^2} + O(t^{-3})$$

where $E(\lambda, Y)$ denotes the extremal length of λ on the Riemann surface Y . The implicit constants depend only on $\chi(S)$.

Note that Theorem 6.2 includes the results stated in the introduction as Theorem 1.4. Before giving the proof of Theorem 6.2, we fix notation and recall some concepts from Teichmüller theory used therein.

Annuli. Let A be an annular Riemann surface of modulus M and let $E(A) = 1/M = E(\gamma, A)$ be the extremal length of γ , the nontrivial isotopy class of simple closed curves on A . Then A is isomorphic to a rectangle $R_A = [0, E(A)] \times (0, 1) \subset \mathbb{C}$ with its vertical sides identified. We call the complex local coordinate z on A coming from this realization the *natural coordinate* for A . Similarly, the induced flat metric $|dz|$ on A is the *natural metric*, with respect to which A is a Euclidean cylinder of height 1 and circumference $E(A)$.

Jenkins-Strebel differentials. For any isotopy class γ of simple closed curves on a compact Riemann surface X , there is a unique embedded annulus $A_\gamma \subset X$ homotopic to γ of maximum modulus $\mathrm{Mod}(A_\gamma) = 1/E(\gamma, X)$. The annulus A_γ is dense in Y , and if z is the natural coordinate for A_γ , the quadratic differential dz^2 on A extends holomorphically to a quadratic differential on X , the *Jenkins-Strebel differential* for γ .

Foliations. A holomorphic quadratic differential ϕ on a Riemann surface X has an associated singular *horizontal foliation* $\mathcal{F}(\phi)$ whose leaves integrate the distribution of tangent vectors v satisfying $\phi(v) \geq 0$. Integration of $|\mathrm{Im} \sqrt{\phi}|$ gives a transverse measure on $\mathcal{F}(\phi)$. Similarly $\mathcal{F}(-\phi)$ is the *vertical foliation*, whose transverse measure comes from $|\mathrm{Re} \sqrt{\phi}|$.

When ϕ is a Jenkins-Strebel differential on a compact surface, the nonsingular leaves of $\mathcal{F}(\phi)$ are closed and homotopic to γ ; in the realization of the Jenkins-Strebel annulus A_γ as a rectangle with identifications, these are the

horizontal lines, while leaves of $\mathcal{F}(-\phi)$ are vertical lines. The transverse measures for $\mathcal{F}(\phi)$ and $\mathcal{F}(-\phi)$ are given by $|dy|$ and $|dx|$ in the rectangle, respectively. Thus the closed leaves of $\mathcal{F}(\phi)$ have total measure $E(A_\gamma) = E(\gamma, X)$ with respect to the transverse measure of the vertical foliation $\mathcal{F}(-\phi)$. Furthermore these closed leaves realize the minimum transverse measure among all curves homotopic to γ (see [Gar, Lem. 11.5.3]).

Pairing and extremal length. The natural pairing between Beltrami differentials $\mu = \mu(z)d\bar{z}/dz$ (with $\mu(z) \in L^\infty$) and integrable holomorphic quadratic differentials $\phi = \phi(z)dz^2$ on a Riemann surface X is given by

$$\langle \mu, \phi \rangle = \operatorname{Re} \int_X \mu \phi = \int_X \mu(z) \phi(z) |dz|^2.$$

When $X = A$ is an annulus with natural coordinate z_A , pairing a Beltrami differential with $\phi = dz_A^2$ gives the infinitesimal change in extremal length $E(A)$ (see [Gar, §1.9]); that is, if A_t is a family of annuli identified by a family of quasiconformal maps with derivative μ_t , then

$$(6.1) \quad \frac{d}{dt} E(A_t) = 2 \langle \mu(t), dz_{A_t}^2 \rangle.$$

Similarly, when ϕ is a Jenkins-Strebel differential on a compact surface X , pairing with ϕ gives the differential of the extremal length function on Teichmüller space:

Theorem 6.3 (Gardiner [Gar, Thm. 11.8.5]). *Let $t \mapsto X_t \in \mathcal{T}(S)$ be a differentiable path whose tangent vector is represented by the Beltrami differential $\mu(t)$ on X_t . Let γ be an isotopy class of simple closed curves and $E(t) = E(\gamma, X_t)$ its extremal length on X_t . Then*

$$E'(t) = 2 \langle \mu(t), \phi(t) \rangle$$

where $\phi(t)$ is the Jenkins-Strebel differential for γ on X_t .

Stretching annuli. Let X be a compact Riemann surface and $A \subset X$ an annulus in the homotopy class of γ , a simple closed curve. The natural coordinate z on A gives a Beltrami differential $d\bar{z}/dz$ on A , which extends to a Beltrami differential on X by setting it to zero on $(X - A)$. This differential represents an infinitesimal affine stretch of A . We will be interested in the extent to which $d\bar{z}/dz$ affects the extremal length of γ , as estimated in

Lemma 6.4. *Let $A \subset X$, γ , and z be as above, and let ϕ be the Jenkins-Strebel differential on X for γ . Then*

$$E(X) \geq \left\langle \frac{d\bar{z}}{dz}, \phi \right\rangle \geq \frac{2E(X)^2}{E(A)} - E(X)$$

where $E(A) = E(\gamma, A)$ and $E(X) = E(\gamma, X)$ are the extremal lengths of γ on these two surfaces.

Remarks.

1. When $A = A_\gamma$ is the Jenkins-Strebel annulus (of maximum modulus), we have $E(A) = E(X)$ and both inequalities in Lemma 6.4 become equalities. However this is clear since $d\bar{z}/dz = \bar{\phi}/|\phi|$ if z is the natural coordinate of the Jenkins-Strebel annulus. The point of the Lemma is that we also have tight bounds for the pairing when A has *nearly* maximum modulus.
2. A related estimate for nearly maximal annuli is used in Kerckhoff's proof of that that foliation map $\mathcal{F} : Q(X) \rightarrow \mathcal{ML}(S)$ is a homeomorphism, see [Ker2, Lem. 3.2].

Proof. Throughout the proof we use the natural coordinate z to identify A with a rectangle $R_A \subset \mathbb{C}$ whose vertical sides are identified.

Writing the restriction of ϕ to A in terms of the natural coordinate, we have

$$\phi = \phi(z)dz^2 = (\alpha + i\beta)^2 dz^2,$$

where α and β are real-valued functions defined locally up to a common sign away from the zeros of ϕ ; in particular the functions α^2 , β^2 and $|\alpha|$ are well-defined almost everywhere. We want to estimate

$$\begin{aligned} \left\langle \frac{d\bar{z}}{dz}, \phi \right\rangle &= \operatorname{Re} \int_A \frac{d\bar{z}}{dz} \phi(z) dz^2 \\ &= \int_A \operatorname{Re} ((\alpha + i\beta)^2) |dz|^2 \\ &= \int_0^1 \int_0^{E(A)} (\alpha^2 - \beta^2) dx dy. \end{aligned}$$

Note that the pairing is computed as an integral over A , rather than X , because the Beltrami coefficient $d\bar{z}/dz$ is understood to be zero on $(X - A)$.

We first derive the upper bound on the pairing. Since A is a subset of X , we have

$$(6.2) \quad \int_0^1 \int_0^{E(A)} (\alpha^2 + \beta^2) dx dy = \int_A |\phi| \leq \int_X |\phi| = E(X).$$

Since $\beta^2 \geq 0$, the same upper bound applies to the integral of $\alpha^2 - \beta^2$, giving

$$\left\langle \frac{d\bar{z}}{dz}, \phi \right\rangle = \int_0^1 \int_0^{E(A)} (\alpha^2 - \beta^2) dx dy \leq E(X).$$

To establish the lower bound on the pairing, note that the horizontal lines in R_A represent closed curves in X homotopic to γ , so the total transverse measure of any one of these with respect to $\mathcal{F}(-\phi)$ is at least $E(X)$. The transverse measure of a curve is the integral of $|\operatorname{Re} \sqrt{\phi}| = |\alpha dx - \beta dy|$, but $dy = 0$ on horizontal lines, so we have

$$\int_0^{E(A)} |\alpha(x + iy)| dx \geq E(X)$$

for all $y \in [0, 1]$. Integrating over y and applying the Cauchy-Schwarz inequality, we obtain

$$(6.3) \quad \int_0^1 \int_0^{E(A)} \alpha^2 dx dy \geq \frac{E(X)^2}{E(A)}.$$

Multiplying (6.3) by 2 and subtracting (6.2) we have

$$\left\langle \frac{d\bar{z}}{dz}, \phi \right\rangle = \int_0^1 \int_0^{E(A)} (\alpha^2 - \beta^2) dx dy \geq \frac{2E(X)^2}{E(A)} - E(X).$$

□

Note that the proof of Lemma 6.4 is essentially a calculation on the annulus A and uses little about the enclosing surface X except that it is foliated by closed trajectories of ϕ . Indeed, the same argument can be applied with the compact surface X replaced by an annulus B and with $\phi = dz_B^2$ the natural quadratic differential on B , and it is this version we will need in the proof of Theorem 6.6:

Lemma 6.5. *Let B be an annular Riemann surface of finite modulus and $A \subset B$ a homotopically essential subannulus. Then*

$$E(B) \geq \left\langle \frac{d\bar{z}_A}{dz_A}, dz_B^2 \right\rangle \geq \frac{2E(B)^2}{E(A)} - E(B)$$

Extremal length and grafting rays. Using Lemma 6.4 as the main technical tool, we are now ready to give the

Proof of Theorem 6.2. First, we consider the case when $\lambda = \gamma$ is a simple closed geodesic with unit weight. For brevity let $Y_t = \text{gr}_{t\gamma} X$; we abbreviate $E(t) = E(\gamma, Y_t)$ and $\ell = \ell(\gamma, X)$.

The proof of (1) follows the usual pattern for an extremal length estimate (see [Ker2, §3]): a particular annulus homotopic to γ bounds $E(t)$ from above, while a particular conformal metric on the surface bounds $E(t)$ from below. In this case the annulus is the grafting cylinder $A_t \subset Y_t$ of modulus t/ℓ , and the conformal metric on Y_t is the *Thurston metric*—the union of the product metric on $A_t = [0, t] \times \gamma$ and the hyperbolic metric of X (see [Tan, §2.1]). Applying the geometric and analytic definitions of extremal length gives

$$\frac{\ell}{t} > E(\gamma, Y_t) > \frac{\ell^2}{t\ell + A} > \frac{\ell}{t} - \frac{A}{t^2}$$

where $A = 4\pi(g-1)$ and $(t\ell + A)$ is the area of the Thurston metric; thus (1) follows.

For statement (2), we must estimate the derivative of extremal length. For all $s, t > 0$, there is a natural quasiconformal map from Y_t to Y_s that is affine on A_t , stretching it vertically in the natural coordinate, and conformal

on $(Y_t - A_t)$. The derivative of this family of maps at $s = t$ is the Beltrami differential

$$\mu(t) = -(2t)^{-1} \frac{d\bar{z}_t}{dz_t}$$

where z_t is the natural coordinate on the grafting annulus A_t and the Beltrami differential $d\bar{z}_t/dz_t$ is understood to be identically zero outside A_t .

By Theorem 6.3, the derivative of extremal length along the grafting ray is

$$(6.4) \quad \frac{d}{dt} E(t) = 2\langle \mu(t), \phi(t) \rangle = -t^{-1} \left\langle \frac{d\bar{z}_t}{dz_t}, \phi(t) \right\rangle.$$

We estimate the pairing $\langle d\bar{z}_t/dz_t, \phi(t) \rangle$ using Lemma 6.4; starting with the upper bound, we have

$$(6.5) \quad \left\langle \frac{d\bar{z}_t}{dz_t}, \phi(t) \right\rangle \leq E(t) \leq \frac{\ell}{t}$$

while the lower bound from the lemma gives

$$\left\langle \frac{d\bar{z}_t}{dz_t}, \phi(t) \right\rangle \geq \frac{2E(t)^2}{E(A_t)} - E(t) = \frac{2t}{\ell} E(t)^2 - E(t).$$

Using $E(t) \geq \ell/t - A/t^2$ on the first term and $E(t) \leq \ell/t$ on the second, we obtain

$$(6.6) \quad \left\langle \frac{d\bar{z}_t}{dz_t}, \phi(t) \right\rangle \geq \frac{2t}{\ell} \left(\frac{\ell}{t} - \frac{A}{t^2} \right)^2 - \frac{\ell}{t} \geq \frac{\ell}{t} - \frac{4A}{t^2}.$$

Multiplying (6.5) and (6.6) by $-1/t$ and using the formula (6.4) for the derivative of extremal length, we have

$$-\frac{\ell(\gamma, X)}{t^2} \leq \frac{d}{dt} E(\gamma, \text{gr}_{t\gamma} X) \leq -\frac{\ell(\gamma, X)}{t^2} + \frac{4A}{t^3},$$

which gives part (2), completing the proof of Theorem 6.2 for simple closed geodesics.

The limiting arguments that extend (1) and (2) to general laminations are completely analogous, so we will only give details for the former. Given a measured lamination $\lambda \in \mathcal{ML}(S)$, define

$$\delta(s, \lambda, X) = s^2 \left| E(\lambda, \text{gr}_{s\lambda} X) - \frac{\ell(\lambda, \text{gr}_{s\lambda} X)}{s} \right|.$$

Since hyperbolic length, extremal length, and grafting are continuous on $\mathcal{ML}(S) \times \mathcal{T}(S)$, this is a continuous nonnegative function $\delta : \mathbb{R}^+ \times \mathcal{ML}(S) \times \mathcal{T}(S) \rightarrow \mathbb{R}$. Part (1) of the Theorem is equivalent to the statement that δ is bounded. We have shown that this is true for simple closed geodesics, i.e. there exists $C > 0$ depending only on $\chi(S)$ such that for all simple closed curves γ and all $s > 0$, we have

$$\delta(s, \gamma, X) \leq C.$$

Suppose that $\lambda = c\gamma$ is a weighted simple closed geodesic. Since hyperbolic length scales linearly in the transverse measure, while extremal length scales quadratically, we have

$$\begin{aligned}\delta(s, c\gamma, X) &= s^2 \left| E(c\gamma, \text{gr}_{sc\gamma} X) - \frac{\ell(c\gamma, \text{gr}_{sc\gamma} X)}{s} \right| \\ &= s^2 c^2 \left| E(\gamma, \text{gr}_{sc\gamma} X) - \frac{\ell(\gamma, \text{gr}_{sc\gamma} X)}{sc} \right| \\ &= \delta(sc, \gamma, X) \leq C.\end{aligned}$$

Since weighted simple closed geodesics are dense in $\mathcal{ML}(S)$, and δ is continuous, this shows that $\delta \leq C$, establishing (1) for all measured laminations.

A similar limiting argument applies to part (2), where one bounds the error function

$$\Delta(s, \lambda, X) = s^3 \left| \left(\frac{d}{ds} E(\lambda, \text{gr}_{s\lambda} X) \right) + \frac{\ell(\lambda, X)}{s^2} \right|.$$

In this case, the function Δ is continuous because extremal length is C^1 on Teichmüller space (see [GM]), with its derivative varying continuously in $\mathcal{ML}(S)$, and since the derivatives of grafting rays depend continuously on the lamination (by Theorem 6.1). $\square$

As a consequence of Theorem 6.2, extremal length decreases along grafting rays outside of a compact set in $\mathcal{T}(S)$ of the form $\{\text{gr}_\lambda X \mid \ell(\lambda, X) \leq L\}$.

Hyperbolic length. The hyperbolic length of λ along a grafting ray is more difficult to control than the extremal length, but for the case of a single curve with large weight, the same techniques used in the proof of Theorem 6.2 give:

Theorem 6.6. *Let $X \in \mathcal{T}(S)$ and let γ be a simple closed hyperbolic geodesic on X . Then the hyperbolic length of γ is of order $1/t$ on the γ -grafting ray and is monotone decreasing for all $t \gg 0$. Specifically, we have*

$$\begin{aligned}(1) \quad \ell(\gamma, \text{gr}_{t\gamma} X) &= \frac{\pi \ell(\gamma, X)}{t} + O(t^{-2}) \\ (2) \quad \frac{d}{dt} \ell(\gamma, \text{gr}_{t\gamma} X) &= -\frac{\pi \ell(\gamma, X)}{t^2} + O(t^{-3})\end{aligned}$$

as $t \rightarrow \infty$, where the implicit constants depend on X and γ .

Note that Theorem 6.6 includes the results stated in the introduction as Theorem 1.5.

Proof. As before let $Y_t = \text{gr}_{t\gamma} X$, and abbreviate $\ell(t) = \ell(\gamma, Y_t)$ and $E(t) = E(\gamma, Y_t)$.

A standard argument using the collar lemma shows that the extremal and hyperbolic length of a curve are asymptotically proportional when the length is small; specifically, we have

$$(6.7) \quad \pi E(t) - CE(t)^2 < \ell(t) < \pi E(t)$$




FIGURE 4. The grafting annulus and its lifts to the annular cover.

for all t such that $E(t) < 1$, where C is a universal constant. Since $E(t) < 1$ for all $t > \ell(\gamma, X)$, substituting the estimate for $E(t)$ from Theorem 6.2 gives (1).

Now we establish the derivative estimate (2). Let $\hat{Y}_t$ denote the cover of Y_t corresponding to the subgroup $\langle \gamma \rangle \subset \pi_1(Y_t)$. Since $\hat{Y}_t$ is conformally equivalent to an annulus of modulus $\pi/\ell(t)$, we have $\ell'(t) = \pi \frac{d}{dt} E(\hat{Y}_t)$.

Recall that $\mu(t) = (-2t)^{-1} d\bar{z}_t/dz_t$ represents the derivative of Y_t , where z_t is the natural coordinate on A_t . Thus the derivative of the annular covers $\hat{Y}_t$ is represented by the pullback Beltrami differential $p^*(\mu(t))$ where $p : \hat{Y}_t \rightarrow Y_t$ is the covering projection. Let w_t denote the natural coordinate on $\hat{Y}_t$. By (6.1) we have

$$(6.8) \quad \ell'(t) = \pi \frac{d}{dt} E(\hat{Y}_t) = 2\pi \langle p^*(\mu(t)), dw_t^2 \rangle.$$

To estimate this pairing, we analyze the differential $p^*(\mu(t))$; its support is the preimage of the grafting cylinder $p^{-1}(A_t) \subset \hat{Y}_t$, which consists of

- (i) A homotopically essential annulus $\hat{A}_t$ such that $p|_{\hat{A}_t} : \hat{A}_t \rightarrow A_t$ is a conformal isomorphism, and
- (ii) A complementary set $\Omega = p^{-1}(A_t) - \hat{A}_t$ that is a disjoint union of countably many simply connected regions $\Sigma_i \subset \hat{Y}_t$ such that the restriction of p to any one of them gives a universal covering $p|_{\Sigma_i} : \Sigma_i \rightarrow A_t$,

as depicted in Figure 4. Therefore we have

$$(6.9) \quad \langle p^*(\mu(t)), dw_t^2 \rangle = \int_{\hat{A}_t} p^*(\mu(t)) dw_t^2 + \int_{\Omega} p^*(\mu(t)) dw_t^2,$$

and we can analyze these two terms individually.

By the length estimates of part (1), the reciprocal moduli $E(\hat{A}_t) = E(A_t) = \ell(0)/t$ and $E(\hat{Y}_t) = \ell(t)/\pi$ differ by $O(t^{-2})$. It follows that the subannulus $\hat{A}_t$ accounts for nearly all of the area of $\hat{Y}_t$. In fact, restricting the metric $|dw_t|$ to $\hat{A}_t$ and using the analytic definition of extremal length gives

$$\text{Area}(\hat{A}_t, |dw_t|) \geq \frac{\ell(\gamma, |dw_t|)^2}{E(\hat{A}_t)} = \frac{E(\hat{Y}_t)^2}{E(\hat{A}_t)} \geq \frac{\ell(0)}{t} - \frac{C}{t^2},$$

where C depends on γ and X . Since $\text{Area}(\hat{Y}_t, |dw_t|) = \ell(0)/t$ and Ω is disjoint from $\hat{A}_t$, we have $\text{Area}(\Omega, |dw_t|) \leq C/t^2$.

This area estimate implies that the second term in (6.9) is negligible, i.e.

$$\begin{aligned} \left| \int_{\Omega} p^*(\mu(t)) dw_t^2 \right| &\leq \|p^*(\mu(t))\|_{\infty} \text{Area}(\Omega, |dw_t|) \\ (6.10) \quad &\leq \left(\frac{1}{2t} \right) \left(\frac{C_2}{t^2} \right) = O(t^{-3}). \end{aligned}$$

Now we consider the first term in (6.9). Since $p|_{\hat{A}_t}$ is a conformal isomorphism, we have $p^*(\mu(t))|_{\hat{A}_t} = (-2t)^{-1} d\bar{z}_t/dz_t$ where $\hat{z}_t$ is the natural coordinate of $\hat{A}_t$. Applying Lemma 6.5 we have

$$E(\hat{Y}_t) \geq \left\langle \frac{d\bar{z}_t}{dz_t}, dw_t^2 \right\rangle \geq \frac{2E(\hat{Y}_t)^2}{E(\hat{A}_t)} - E(\hat{Y}_t).$$

As before we substitute $E(\hat{Y}_t) = \ell(t)/\pi$, $E(\hat{A}_t) = \ell(0)/t$, and apply the estimates for $\ell(t)$ to obtain

$$\int_{\hat{A}_t} p^*(\mu(t)) dw_t^2 = (-2t)^{-1} \left\langle \frac{d\bar{z}_t}{dz_t}, dw_t^2 \right\rangle = -\frac{\ell(0)}{2t^2} + O(t^{-3}).$$

Thus we have estimates for both terms in (6.9), and applying the formula (6.8) for $\ell'(t)$ gives the desired result:

$$\ell'(t) = -\frac{\pi\ell(0)}{t^2} + O(t^{-3}).$$

□

Remark. In [McM, Cor. 3.2], McMullen shows that $\ell(\lambda, \text{gr}_{t\lambda} X) < \ell(\lambda, X)$ for all $X \in \mathcal{T}(S)$, $\lambda \in \mathcal{ML}(S)$ and $t > 0$; it is also mentioned that this upper bound can be strengthened to

$$\ell(\gamma, \text{gr}_{t\gamma} X) \leq \frac{\pi}{\pi + t} \ell(\gamma, X),$$

for any simple closed curve γ . For details on this and a corresponding lower bound, see [DK, Prop. 3.4]. Theorem 6.6 shows that this upper bound is asymptotically sharp.

REFERENCES

- [Ber1] Lipman Bers. Simultaneous uniformization. *Bull. Amer. Math. Soc.* **66**(1960), 94–97.
- [Ber2] Lipman Bers. Correction to “Spaces of Riemann surfaces as bounded domains”. *Bull. Amer. Math. Soc.* **67**(1961), 465–466.
- [Bon1] Francis Bonahon. Earthquakes on Riemann surfaces and on measured geodesic laminations. *Trans. Amer. Math. Soc.* **330**(1992), 69–95.
- [Bon2] Francis Bonahon. Shearing hyperbolic surfaces, bending pleated surfaces and Thurston’s symplectic form. *Ann. Fac. Sci. Toulouse Math. (6)* **5**(1996), 233–297.
- [Bon3] Francis Bonahon. Geodesic laminations with transverse Hölder distributions. *Ann. Sci. École Norm. Sup. (4)* **30**(1997), 205–240.
- [Bon4] Francis Bonahon. Transverse Hölder distributions for geodesic laminations. *Topology* **36**(1997), 103–122.
- [Bon5] Francis Bonahon. Variations of the boundary geometry of 3-dimensional hyperbolic convex cores. *J. Differential Geom.* **50**(1998), 1–24.
- [Bon6] Francis Bonahon. Kleinian groups which are almost Fuchsian. *J. Reine Angew. Math.* **587**(2005), 1–15.
- [BO] Francis Bonahon and Jean-Pierre Otal. Laminations mesurées de plissage des variétés hyperboliques de dimension 3. *Ann. of Math. (2)* **160**(2004), 1013–1055.
- [DK] Raquel Díaz and Inkang Kim. Asymptotic Behavior of Grafting Rays. Preprint, 2007.
- [D1] David Dumas. Grafting, pruning, and the antipodal map on measured laminations. *J. Differential Geometry* **74**(2006), 93–118. Erratum. *J. Differential Geometry* **77**(2007), 175–176.
- [D2] David Dumas. The Schwarzian derivative and measured laminations on Riemann surfaces. *Duke Math. J.* **140**(2007), 203–243.
- [EM] D. B. A. Epstein and A. Marden. Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces. In *Analytical and geometric aspects of hyperbolic space (Coventry/Durham, 1984)*, volume 111 of *London Math. Soc. Lecture Note Ser.*, pages 113–253. Cambridge Univ. Press, Cambridge, 1987.
- [EMM] D. B. A. Epstein, A. Marden, and V. Markovic. Quasiconformal homeomorphisms and the convex hull boundary. *Ann. of Math. (2)* **159**(2004), 305–336.
- [GKM] Daniel Gallo, Michael Kapovich, and Albert Marden. The monodromy groups of Schwarzian equations on closed Riemann surfaces. *Ann. of Math. (2)* **151**(2000), 625–704.
- [Gar] Frederick P. Gardiner. *Teichmüller theory and quadratic differentials*. Pure and Applied Mathematics (New York). John Wiley & Sons Inc., New York, 1987. , A Wiley-Interscience Publication.
- [GM] Frederick P. Gardiner and Howard Masur. Extremal length geometry of Teichmüller space. *Complex Variables Theory Appl.* **16**(1991), 209–237.
- [Gol] William M. Goldman. Projective structures with Fuchsian holonomy. *J. Differential Geom.* **25**(1987), 297–326.
- [Hej] Dennis A. Hejhal. Monodromy groups and linearly polymorphic functions. *Acta Math.* **135**(1975), 1–55.
- [KT] Yoshinobu Kamishima and Ser P. Tan. Deformation spaces on geometric structures. In *Aspects of low-dimensional manifolds*, volume 20 of *Adv. Stud. Pure Math.*, pages 263–299. Kinokuniya, Tokyo, 1992.
- [Kat] A. B. Katok. Invariant measures of flows on orientable surfaces. *Dokl. Akad. Nauk SSSR* **211**(1973), 775–778.
- [Ker1] S. P. Kerckhoff. Simplicial systems for interval exchange maps and measured foliations. *Ergodic Theory Dynam. Systems* **5**(1985), 257–271.

- [Ker2] Steven P. Kerckhoff. The asymptotic geometry of Teichmüller space. *Topology* **19**(1980), 23–41.
- [Ker3] Steven P. Kerckhoff. Earthquakes are analytic. *Comment. Math. Helv.* **60**(1985), 17–30.
- [Kle] F. Klein. *Vorlesungen Über die Hypergeometrische Funktion*. Springer-Verlag, Berlin, 1933.
- [Lev] Gilbert Levitt. Feuilletages des surfaces. *Ann. Inst. Fourier (Grenoble)* **32**(1982), x, 179–217.
- [Mas1] Bernard Maskit. On a class of Kleinian groups. *Ann. Acad. Sci. Fenn. Ser. A I No.* **442**(1969), 8.
- [Mas2] Howard Masur. Interval exchange transformations and measured foliations. *Ann. of Math. (2)* **115**(1982), 169–200.
- [McM] Curtis T. McMullen. Complex earthquakes and Teichmüller theory. *J. Amer. Math. Soc.* **11**(1998), 283–320.
- [Nag] Subhashis Nag. *The complex analytic theory of Teichmüller spaces*. Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons Inc., New York, 1988. A Wiley-Interscience Publication.
- [Ota] Jean-Pierre Otal. Le théorème d’hyperbolisation pour les variétés fibrées de dimension 3. *Astérisque* **235**(1996), x+159.
- [Pap] Athanase Papadopoulos. Deux remarques sur la géométrie symplectique de l’espace des feuilletages mesurés sur une surface. *Ann. Inst. Fourier (Grenoble)* **36**(1986), 127–141.
- [PH] R. C. Penner and J. L. Harer. *Combinatorics of train tracks*, volume 125 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 1992.
- [Pla] Ioannis D. Platis. Complex symplectic geometry of quasi-Fuchsian space. *Geom. Dedicata* **87**(2001), 17–34.
- [Ree] Mary Rees. An alternative approach to the ergodic theory of measured foliations on surfaces. *Ergodic Theory Dynamical Systems* **1**(1981), 461–488 (1982).
- [SW] Kevin P. Scannell and Michael Wolf. The grafting map of Teichmüller space. *J. Amer. Math. Soc.* **15**(2002), 893–927 (electronic).
- [Ser1] Caroline Series. On Kerckhoff minima and pleating loci for quasi-Fuchsian groups. *Geom. Dedicata* **88**(2001), 211–237.
- [Ser2] Caroline Series. Limits of quasi-Fuchsian groups with small bending. *Duke Math. J.* **128**(2005), 285–329.
- [Ser3] Caroline Series. Thurston’s bending measure conjecture for once punctured torus groups. In *Spaces of Kleinian groups*, volume 329 of *London Math. Soc. Lecture Note Ser.*, pages 75–89. Cambridge Univ. Press, Cambridge, 2006.
- [SB] Yaşar Sözen and Francis Bonahon. The Weil-Petersson and Thurston symplectic forms. *Duke Math. J.* **108**(2001), 581–597.
- [ST] Dennis Sullivan and William Thurston. Manifolds with canonical coordinate charts: some examples. *Enseign. Math. (2)* **29**(1983), 15–25.
- [Tan] Harumi Tanigawa. Grafting, harmonic maps and projective structures on surfaces. *J. Differential Geom.* **47**(1997), 399–419.
- [Thu1] William P. Thurston. Geometry and Topology of Three-Manifolds. Princeton lecture notes, 1979. <http://www.msri.org/publications/books/gt3m/>
- [Thu2] William P. Thurston. Earthquakes in two-dimensional hyperbolic geometry. In *Low-dimensional topology and Kleinian groups (Coventry/Durham, 1984)*, volume 112 of *London Math. Soc. Lecture Note Ser.*, pages 91–112. Cambridge Univ. Press, Cambridge, 1986.
- [Thu3] William P. Thurston. Minimal stretch maps between hyperbolic surfaces. Unpublished preprint, 1986. [arXiv:math.GT/9801039](https://arxiv.org/abs/math.GT/9801039)

- [Tro] Anthony J. Tromba. *Teichmüller theory in Riemannian geometry*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 1992. Lecture notes prepared by Jochen Denzler.
- [Vee] William A. Veech. Gauss measures for transformations on the space of interval exchange maps. *Ann. of Math. (2)* **115**(1982), 201–242.
- [Wol1] Scott Wolpert. An elementary formula for the Fenchel-Nielsen twist. *Comment. Math. Helv.* **56**(1981), 132–135.
- [Wol2] Scott Wolpert. The Fenchel-Nielsen deformation. *Ann. of Math. (2)* **115**(1982), 501–528.
- [Wol3] Scott Wolpert. On the symplectic geometry of deformations of a hyperbolic surface. *Ann. of Math. (2)* **117**(1983), 207–234.
- [Wol4] Scott A. Wolpert. Chern forms and the Riemann tensor for the moduli space of curves. *Invent. Math.* **85**(1986), 119–145.
- [ZT] P. G. Zograf and L. A. Takhtajan. On the Liouville equation, accessory parameters and the geometry of Teichmüller space for Riemann surfaces of genus 0. *Mat. Sb. (N.S.)* **132(174)**(1987), 147–166.

BROWN UNIVERSITY

E-mail address: ddumas@math.brown.edu

URL: <http://www.math.brown.edu/~ddumas/>

RICE UNIVERSITY

E-mail address: mwolf@math.rice.edu

URL: <http://www.math.rice.edu/~mwolf/>